



# Geological factors affecting the chemical characteristics of the thermal waters of the carbonate karstified aquifers of Northern Vietnam

C. Drogue, N. N. Cat, J. Dazy

## ► To cite this version:

C. Drogue, N. N. Cat, J. Dazy. Geological factors affecting the chemical characteristics of the thermal waters of the carbonate karstified aquifers of Northern Vietnam. Hydrology and Earth System Sciences Discussions, 2000, 4 (2), pp.332-340. 10.5194/hess-4-332-2000 . hal-00330836

**HAL Id: hal-00330836**

**<https://hal.science/hal-00330836>**

Submitted on 1 Jan 2000

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Geological factors affecting the chemical characteristics of the thermal waters of the carbonate karstified aquifers of Northern Vietnam

C. Drogue<sup>1</sup>, N. N. Cat<sup>2</sup> and J. Dazy<sup>1</sup>

<sup>1</sup>Hydrogéologie, UMR 5569 Hydrosiences, Université Montpellier 2, place Eugène Bataillon, 34095 Montpellier Cedex 5, France

<sup>2</sup>Institute of Geography, Vietnam National Centre for Sciences and Technology, Nghia Do, Tu Liem, Hanoi, Vietnam

e-mail for corresponding author: drogue@dstu.univ-montp2.fr

## Abstract

In northern Vietnam, exposed carbonate rock formations cover an area of more than 50,000 km<sup>2</sup>. Their accumulated thickness from the Cambrian to the Triassic is in some places as much as 3000 m. Numerous thermal waters (springs and wells) occur in these strongly karstified carbonate massifs. This is the result of significant ancient and present orogenic activity, as the region demonstrates by its strong seismic activity. These karstic formations are water-bearing and strongly recharged by rainfall of between 1600 mm and 2000 mm per year in 90% of the area concerned. In view of the average annual air temperatures (17°C–25°C according to the region), 23 sample springs or wells were chosen with water temperatures of between 29°C and 68°C. Hydrochemical characteristics of these thermal waters emerging in different carbonate-rock units were examined by chemical analyses of major ions. In this large region, thermal waters are divided into four hydrochemical types: the Na-Cl type resulting from the intrusion of sea water for distances of up to several kilometres inland and depths of 1000 m, the Ca-SO<sub>4</sub> type, probably resulting from the leaching of deposits of metallic sulphides that are widely distributed in these carbonate-rock units, and finally the Ca-HCO<sub>3</sub> and Mg-HCO<sub>3</sub> types which are chemically similar to fresh karstic waters in limestones and dolostones. The occurrence of these thermal groundwaters as well as their chemical characteristics seem to indicate the existence of large-scale deepseated groundwater flow systems in the karstic aquifers.

**Keywords:** Vietnam; thermal waters; karst; hydrochemistry

## Introduction

South-East Asia is rich in thermal springs (Grimaud *et al.*, 1985; Shen Zhian *et al.*, 1989) which may flow from either eruptive or volcanic rocks or from sedimentary rocks. In the latter case, the occurrence of thermal water in carbonate rocks is of particular interest. Indeed, these rocks are often karstified, thus forming groundwater reservoirs in which flow resulting from the infiltration of rain water can be substantial. The presence of hot water means that deep thermal flows are sufficiently active not to be masked wholly at emergence by mixing with cold water. It is, therefore, possible that the hydrochemical characteristics may provide information about the modes of deep flow and, in particular, may show whether the flowpath of groundwater is related to formations other than carbonate rocks.

Northern Vietnam (north of the 17th parallel) possesses large exposed units of carbonate rocks covering a sixth of the territory, an area of nearly 50,000 km<sup>2</sup> (Fig. 1). These carbonate rocks are karstified and feed a large number of springs. They offer a resource which is exploited by wells for water supply. The water temperatures are abnormally

high in some places. The intention was to identify the major chemical characteristics in a preliminary study of the karstic thermal waters. This could demonstrate the hydrochemical homogeneity or diversity at the scale of the region and the possibilities of deep flows from non-carbonate rocks.

## The geological and climatic frameworks

In northern Vietnam, the oldest carbonate formations (Proterozoic and Lower Cambrian) are represented by marbles. However, the main carbonate rocks in the region, consisting of marble, limestone and dolomite were formed at the middle Cambrian, Ordovician, Devonian, Carboniferous and Permian periods. The cumulated thickness of these carbonate formations can reach 3,000 m (Fromaget, 1941).

Northern Vietnam is an area of substantial tectonic movement due to the presence of one of the main active strike-slip faults in South-East Asia, the Red River Fault zone (Fig. 1) (Rangin *et al.*, 1995; Harrison *et al.*, 1996).




Fig. 1. Carbonate rock outcrops in northern Vietnam (after Geological Map of Vietnam, 1986 and unpublished map of CNST-Vietnam). Location of the springs and wells studied.

1. Observed and assumed geological boundary.
2. Observed and assumed fault.
3. Carbonate rock massifs.
4. Submerged karst in Halongbay.
5. Spring/well (the number refers to Tables 1 and 2)
6. Na-Cl typewaters,
7. Ca-SO<sub>4</sub> typewaters,
8. Ca-HCO<sub>3</sub> typewaters,
9. Non thermal waters.

This is the extension towards the south-east of the major fault separating the Indo-Australian plate from the Eurasian plate.

In addition, various tectonic phases (Huang Chu-Ching, 1978; Allen *et al.*, 1984; Leloup *et al.*, 1995) have enhanced the formation of the Red River basin in the Hanoi region with more than 3000 m of sedimentary deposits in the deepest part (Pham *et al.*, 1995). In this alluvial basin, carbonate-rock formations lie at great depth whereas karstic carbonate units outcrop widely in the northern and southern parts of the basin. The appearance of many granitic and rhyolitic intrusions (during the Jurassic and the Cretaceous) are also possible sources of thermal anomalies in the subsurface.

Tectonic movements along the major faults and the related fracturing must enhance the deep flow paths of fluids, especially as the region is the site of strong neotectonic activity. As many as 400 earthquakes have been observed annually (Can and Hoe, 1969; Xuyen, 1991). That of 1 November 1935 scored 9 on the Richter scale and caused the opening of a 50 m long crevasse and gushes of hot water. The hypocentres of the earthquakes in 1956 and

1957 were located on the Red River fault, which is thus currently active (Yem, 1986).

The intense fracturing resulting from these tectonic movements allows the infiltration of water and has enhanced karstification in the carbonate sequences of the geological series. Outcrops of the karstified sequences display different morphological forms including the spectacular tower karst and pinnacle landscape (Blondel, 1929; Glazek, 1966; Fenart *et al.*, 1999). Similar karstification has been observed at depth throughout northern Vietnam in a large number of wells drilled to extract water at depths of up to 1,000 m (wells 18 and 19 listed in Table 1).

The groundwater recharge of these aquifers depends on the climate. According to Tran Cau (1984), northern Vietnam has three types of climate, a humid tropical climate in the low-lying plains (the Red River delta), a highland subtropical climate at elevations of 600 to 2,500 m and a temperate climate in the mountains above 2,500 m. The average annual rainfall is between 1,600 and 2,000 mm in 90% of the region (Fig. 2). In the north-west hills, the rainfall gradients are steep and the precipitation exceeds

**Table 1.** Results of the chemical analyses of the thermal waters. The concentrations are expressed in mg l<sup>-1</sup>.  $\Sigma r^+$  and  $\Sigma r^-$ : Total cations and total anions (expressed in meq/l): Geology, D: Devonian, C: Carboniferous, P: Permian, T: Trias, a ? indicates uncertain determinations. The locations of the sites are indicated on the map in Fig. 1.

N°	Location	T°C	TDS (mgL <sup>-1</sup> )	Ca <sup>++</sup> (mgL <sup>-1</sup> )	Mg <sup>++</sup> (mgL <sup>-1</sup> )	Na <sup>+</sup> (mgL <sup>-1</sup> )	$\Sigma r^+$ (meq/L)	$\Sigma r^-$ (meq/L)	HCO <sub>3</sub> <sup>-</sup> (mgL <sup>-1</sup> )	Cl <sup>-</sup> (mgL <sup>-1</sup> )	SO <sub>4</sub> <sup>--</sup> (mgL <sup>-1</sup> )	Geol
<b>Na-Cl type waters</b>												
2	Tam Hop	42	15,941.5	605.50	390.0	4,868.1	274.00	275.50	137.2	8,842.1	1098.6	C.P.
3	Quang Hanh	42	26,602.4	695.90	812.5	8,104.0	454.00	462.50	211.5	14,851.3	1927.2	C.P.
6	Muong Pia	55	5,083.0	491.00	213.0	916.9	81.90	82.30	338.0	1,588.1	1536.0	?
18	Thach Khôi	44	488.3	45.13	8.7	120.7	8.22	8.01	47.6	231.0	35.2	C
19	Cam Giang	38	1,122.8	128.80	28.3	217.7	18.20	19.10	49.4	509.4	189.2	C
20	Kênh Gà	52	8,871.2	201.80	30.4	3,206.8	151.90	152.00	91.5	5,336.7	4.0	T
22	Tiên Lang	54	13,145.3	566.40	321.4	3,938.1	226.00	226.60	118.5	7,301.3	899.6	T
<b>Ca-SO<sub>4</sub> type waters</b>												
1	Vang Po	37	1,002.0	200.4	60.8	14.0	15.6	14.2	219.7	7.1	500.0	T
5	Phu Mao	40	1,078.2	227.4	36.3	34.3	15.7	15.4	201.3	5.2	573.7	?
8	Ban Peo	47	2,296.4	400.8	109.4	111.7	33.8	33.9	253.7	28.8	1392.0	?
9	Ban Bon	45	2,444.2	501.2	94.7	72.0	36.3	35.7	308.1	13.6	1454.6	?
10	Nong Da	29	2,259.6	538.1	82.2	9.0	34.0	33.1	195.2	5.1	1430.0	?
11	Ban Hóc	53	2,317.8	517.6	79.7	52.0	35.1	33.7	244.1	6.8	1417.6	?
12	Pê Ngoài	38	2,201.5	476.5	104.7	17.0	33.2	32.4	219.7	5.1	1378.5	?
13	Ba Vi	33	1,282.3	292.6	45.0	7.9	18.8	18.7	183.1	7.8	745.9	T
17	Suoi Am	45	1,143.3	228.5	38.4	3.6	14.7	17.2	231.9	7.2	633.7	T
23	Thanh Tan	68	921.6	173.7	20.4	89.2	14.5	12.5	238.5	25.9	373.9	D
<b>Ca-HCO<sub>3</sub> and Mg-HCO<sub>3</sub> type waters</b>												
4	La Hien	33	272.4	43.8	9.7	12.2	3.50	3.51	191.8	7.7	7.2	D
7	Ban Mong	39	420.5	68.1	13.4	18.7	5.36	5.40	311.2	7.1	2.0	T
14	Mo Da	34	427.9	82.6	15.2	4.7	5.62	5.85	232.5	11.7	81.2	T
15	Dam Thi	34	439.5	85.6	12.2	12.9	5.85	5.83	237.9	4.2	86.4	T
16	Qui Hoa	37	341.3	38.0	9.7	30.5	4.00	4.55	220.9	7.1	35.1	T
21	Thuong Xung	37	429.5	18.6	55.1	11.0	5.92	5.77	325.1	5.1	14.6	T
<b>Sea-water</b>			34,430.0	410.0	1,284.0	10,551.0			41.0	19,046.0	2,653.0	

4,000 mm per year at the highest summits (the highest elevation is 3,142 m in the Fan Si Pan range). As 54% of the rainfall does not evaporate (Tran Cau, 1984), the karstic aquifers benefit from ample supplies and flows are thus very active.

Average annual air temperatures are between 22°C and 26°C in the Red River delta and 15°C to 16°C in the mountains. As a result, in this study the underground water whose temperature exceeds 26°C is considered to be thermal water, for all regions.

## The chemical characteristics of the thermal waters

Thermal water is defined here as water whose temperature is higher than the average mean annual temperature on

discharge at the surface. In northern Vietnam, the waters described in this paper as mineral and thermal are spring waters or well waters originating in carbonate karstic rock. Nearly all have at one time been used for medicinal or bottling purposes. Natural flows are between 0.51 s<sup>-1</sup> and 141 s<sup>-1</sup> for springs and between 41 s<sup>-1</sup> and 151 s<sup>-1</sup> for artesian wells.

## SAMPLING AND DATA ANALYSIS

Sampling of deep waters and spring waters from karstic rocks has been carried out recently. The analyses were performed by the University of Mining and Geology in Hanoi and by the Vietnam Institute of Petroleum and Gas using standard methods. The chemical analyses concern only the major ions (Ca, Mg, Na, K and Cl, SO<sub>4</sub>, HCO<sub>3</sub>).




Fig. 2. Average annual rainfall in northern Vietnam. The lines of equal rainfall are in mm. The annual rainfall depth is between 1600 mm and 2000 mm in the hatched zone.

Minor constituents such as bromide, silica ( $\text{SiO}_2$ ), boron and fluoride were assessed in some waters.


Detailed or partial chemical analyses are available for the majority of the 23 thermal and mineral waters examined and a representative selection of these is provided in Table 1. The temperature of these waters ranges from 29°C to 68°C (Fig. 3).

Examination of the ionic balance (concentrations expressed in  $\text{meq l}^{-1}$ ) was used to check the accuracy of the analyses (Standard Methods, 1974). It is considered that deviation of 0.05 in 1, is acceptable in the anion/cation ratio (Domenico and Schwartz, 1990). Here, the deviation is less than  $\pm 0.05$  for 19 analyses. Analyses 1, 16, 17 and 23 for which the deviation is between 0.09 and 0.17 should therefore be suspect.

It will be seen that potassium does not appear in Table 1. Indeed, this element was only analysed in five samples (9, 11, 12, 13 and 14) for which the anion/cation ratio was similar to that of analyses with no value for potassium. The absence of  $\text{K}^+$  does not therefore seem a determinant factor in the value of the anions-cations balance. However, the possibility that the imbalances in analyses 1, 16, 17 and 23 might result partly from the absence of potassium cannot be ruled out.

#### CHEMICAL CHARACTERISTICS OF THE THERMO-MINERAL WATERS

Within the diamond-shaped field of a trilinear diagram (Piper, 1944), four distinct subfields can be identified and the ionic species predominant in each are indicated in Fig. 4.



- △ - Calcium bicarbonate water
- - Calcium sulphate water
- - Sodium chloride water

Fig. 3. Temperature of the thermal waters studied. The hatched zone a-b represents the range of annual average air temperatures for the whole of northern Vietnam.

It is apparent that four types of mineral and thermal waters can be distinguished and the number of waters falling into each group is shown in brackets:

- Na-Cl type (7)
- Ca- $\text{SO}_4$  type (10)
- Ca- $\text{HCO}_3$  type (5)
- Mg- $\text{HCO}_3$  type (1)

Representation on semi-logarithmic diagrams (Schoeller, 1962) makes it possible to compare the hydrochemical types according to the concentrations in  $\text{meq l}^{-1}$ . (Figs. 5 and 8).




Fig. 4. Representation of the chemical analyses of the thermal waters studied (trilinear diagram, after Piper, 1944).

#### THE NA-CL TYPE WATERS (FIG. 5)

The waters belonging to this hydrochemical facies are from five wells (Fig. 5a) and two springs (Fig. 5b).

##### (a) Water from wells (2, 3, 18, 19 and 22)

These wells are in the Haiphong coastal region and located in two regions. Wells 2 and 22 are in karstified limestone of the coast of Halong Bay. Water is extracted at depths of between 100 m and 800 m (22). It is strongly mineralised ( $13.1 < \text{TDS g l}^{-1} < 26$ ) and its temperature ranges from 42°C to 54°C. The chemical composition is characterised by high Na-Cl contents and rNa/Cl and rSO<sub>4</sub>/Cl ratios are close to 0.85 and 0.1 respectively,




Fig. 5. Sodium chloride type waters. (logarithmic diagram, after Schoeller, 1962). 5a: Intrusion of sea water in thermal wells. 5b: Thermo-mineral springs.




Fig. 6. Relation between sodium and chloride in thermal waters. SW = Sea-water, KS = Karstic waters.

identical to sea water rNa/Cl ratio and sea-water rSO<sub>4</sub>/Cl ratio. Chloride and sodium concentrations are plotted in Fig. 6. The relationship observed between Cl<sup>-</sup> and Na<sup>+</sup> close indicates, to a first approximation, a mixing with sea-water.

The use of bromide anion concentrations supplements chloride measurements in explaining the salinity anomalies in ground waters (Morell *et al.*, 1986; Fisher and Mullican, 1997; Andreassen and Fleck, 1997; Hsissou *et al.*, 1999). In sea-water, the Br:Cl ratio is approximately 0.0034. This ratio is 0.0029 for well 2, 0.0036 for well 3, and 0.0024 for well 22 (Table 3). That would confirm the contamination by the sea-water. Wells 18 and 19 tap a karstic aquifer at a depth of between 766 m and 1,017 m under the vast Red River delta alluvial plain. The wells are artesian and the water temperatures are 38°C and 44°C. In wells 18 and 19, the Br:Cl ratios are 0.0104 and 0.0294 respectively. The enrichment in bromide could have been generated by sediments (Neogene and Pleistocene sus-jacents, Fig. 7)

##### (b) Spring water (6 and 20)

Although these two thermal springs also fall on the mixing line (Fig. 6), their geographical situation prevents contamination by sea-water.

The water from Kanh Gà spring (20) emerges at 52°C from a karstic massif lying south of the alluvial plain of the Red River. Its hydrochemical characteristics differ from the Na-Cl type waters analysed in the delta zone. The 8.54 g l<sup>-1</sup> Na-Cl concentration represents 96% of the TDS. The sulphate content (4 mg l<sup>-1</sup>) and the Mg<sup>2+</sup> concentration (30 mg l<sup>-1</sup>) are low in comparison with those of the waters contaminated by sea-water, in which the level is generally more than 300 mg l<sup>-1</sup>.

This depletion of sulphate may be explained by traces of H<sub>2</sub>S observed during sampling. This is commonly accounted for by a bacterial reduction phenomenon; this occurs frequently in the strongly reducing environment of confined aquifers (Schoeller, 1962). The SO<sub>4</sub>/Cl ratio is thus 6.10<sup>-4</sup> here, whereas it is 10<sup>-1</sup> in sea-water.




Fig. 7. Lithologic log of well 18.


The water from Muong Pia spring (6, temperature 55°C) clearly differs from the previous samples. Its Na-Ca-Mg-Cl-SO<sub>4</sub> ionic sequence is unique in the analyses and the rSO<sub>4</sub>/Cl ion ratio (0.7) is higher than that of sea-water. In fact, the geographical position of this spring is 150 km from the coast, ruling out a marine component. This hydro-chemical feature led to the examination of other features. Thus, the concentrations of boron (1.5 mg l<sup>-1</sup>) fluoride (1.6 mg l<sup>-1</sup>) and silica (63 mg l<sup>-1</sup>) may be characteristic of water that has flowed in contact with crystalline formations in the underlying basement.

#### THE CA-SO<sub>4</sub> TYPE WATERS (1, 5, 8, 9, 10, 11, 12, 13, 17 AND 23) (FIG. 8)

There are ten analyses of this type, consisting of nine springs and a well (13) at Ba Vi near Hanoi. The chemical compositions of these sulphate waters can be grouped into three major ionic types: the Ca-SO<sub>4</sub> type where calcium represents more than 20% of the cationic sum (11, 13, 17, 23) (Fig. 8a), the Ca-Mg-SO<sub>4</sub> type (8, 9, 10, 12) and the Ca-Mg-SO<sub>4</sub>-HCO<sub>3</sub> type (1, 5). The temperature of these sulphate waters ranges between 29°C and 68°C. The springs of Ban Hoc (11) and Thanh Tan (23) are, with Muong Pia spring (6–55°C), the warmest (53°C and 68°C) of all the springs in this study.

With TDS of 1 to 2.5 g l<sup>-1</sup> and SO<sub>4</sub><sup>2-</sup> contents of 0.5 to 1.5 g l<sup>-1</sup>, the mineral contents of these waters are much lower than the sodium chloride type waters. Analyses 1, 8, 9, 10 and 12, which display high Mg<sup>2+</sup> content (up to 110 mg l<sup>-1</sup>) belong to the Ca-Mg-SO<sub>4</sub> type (Fig. 8b).

Like the chloride contents (between 5.2 and 28.8 mg l<sup>-1</sup>), the sodium contents were found generally to be low in these

Fig. 8. Calcium sulphate type waters. 8a: Ca-SO<sub>4</sub> type, 8b: Ca-Mg-SO<sub>4</sub> type.

waters with  $3.6 < \text{Na} < 52 \text{ mg l}^{-1}$  in almost all cases. These levels probably indicate inflow of rain water as the source of Na and Cl ions.

For the purpose of the study, sulphate waters can be divided into two groups. The first has a temperature between 45°C and 68°C and includes analyses 8, 9, 11 and 23, which are characterised by relatively high Na<sup>+</sup> levels (72 to 111.7 mg l<sup>-1</sup>). The second group is represented by the waters whose temperature is between 30°C and 45°C and whose Na<sup>+</sup> contents are lower than 35 mg l<sup>-1</sup> (1, 5, 10, 12, 13 and 17). These waters have generally a rNa/Cl ratio ranging between 1 and 5 close to the rNa/Cl ratio of rain waters.

#### THE CA-HCO<sub>3</sub> AND MG-HCO<sub>3</sub> TYPE WATERS (4, 7, 14, 15, 16 AND 21) (FIG. 9A)

These consist of four springs and a well (4, 120 m deep) of the Ca-SO<sub>4</sub><sup>2-</sup> type and a spring of the Mg-HCO<sub>3</sub> type. Water temperatures are between 33°C and 39°C. TDS values are between 272 mg l<sup>-1</sup> and 440 mg l<sup>-1</sup>, indicating the lowest mineralisation of all the waters studied. Spring 21 (Thuong Xung) is the only one of the Mg-HCO<sub>3</sub> type. It flows at the foot of calcareous-dolomitic plateaux (Ninh Binh region). With a content of 55 mg l<sup>-1</sup>, magnesium forms 76% of total cations and the HCO<sub>3</sub><sup>-</sup> content (325 mg l<sup>-1</sup>) is one of the highest after that of Muong Pia (6,338 mg l<sup>-1</sup>). With regard to the other elements, this analysis is similar to that of the Son La karstic cold spring (25), with the same TDS of 430 mg l<sup>-1</sup> and comparable HCO<sub>3</sub><sup>-</sup> level. Only the Mg<sup>2+</sup> content is different, making the rCl/Mg ratio 0.2 rather than 0.7.

## Discussion and conclusion

In all parts of the world, karstic waters are subsurface waters




Fig. 9. Calcium bicarbonate type waters. Examples of (9a) thermal waters and (9b) non-thermal karstic waters.

different from deep-seated thermal waters and they generally display a chemical appearance of the calcium bicarbonate type (Back and Hanshaw, 1965; Hem, 1970). Examples of this water-type in northern Vietnam are found in analyses 24, 25, 26, 27, 28 and 29 (Table 2 and Fig. 9b). In these analyses, mineralisation is similar to that of bicarbonate type thermo-mineral waters (TDS between  $172 \text{ mg l}^{-1}$  and  $485 \text{ mg l}^{-1}$ ). As in most thermal waters, the  $\text{SO}_4^{2-}$  contents are lower than the  $\text{Cl}^-$  contents in all cases. The  $\text{rNa/Cl}$  ratio is close to 1 in the non-thermal waters, which would seem to show the effect of rain containing marine aerosols. In contrast, the values of this ratio differ from 1 in the thermal waters, possibly as a result of ion exchange with clay minerals.

This hydrochemical proximity of these thermal and non-thermal waters indicates that hot water flows only in carbonate rocks but that they are subjected to ionic exchange (Domenico and Schwartz, 1990) as indicated by the  $\text{rNa/Cl}$  ratio.

Calcium sulphate waters are the most frequently seen type. This water type has already been mentioned as being very common in northern Vietnam (Fontaine, 1969). In

Table 3. Bromide concentrations, chloride concentrations and bromide:chloride ratios of the thermo-mineral waters (Na-Cl type). The numbers are indicated on the map in Fig. 1.

N°	Bromide ( $\text{mg L}^{-1}$ )	Chloride ( $\text{mg L}^{-1}$ )	$\text{Br}^-:\text{Cl}^-$ ratio
2	25.6	8,842.1	0.0029
3	53.2	14,851.3	0.0036
18	2.4	231.0	0.0104
19	15.0	509.4	0.0294
22	17.6	7,301.3	0.0024
Sea-water	64.0	19,046.0	0.0034

addition, Waring (1965) reported on several thermal springs in northern Vietnam and noted that the main chemical compound was the  $\text{SO}_4^{2-}$  ion. This type of water may, therefore, be characteristic of the thermal waters in this region. The source of the sulphate cannot originate from dissolution of gypsum or anhydrite as these evaporitic deposits are not found in northern Vietnam. They are reported only in the Cretaceous deposits in central Vietnam (Geological Map of Vietnam, 1986). However, it is known that successive orogenic periods enhanced the magmatic activities that led to much mineralisation in northern Vietnam (Tran Cau, 1984) and especially sulphide mineralisation features (Fromaget, 1941) such as pyrite, molybdenite, arsenopyrite, sphalerite and lead sulphide in the Palaeozoic schists underlying the karst. Permian-Carboniferous limestones also contain pyrite (as in the cores withdrawn from well 13), arsenopyrite and sphalerite and also chalcocite. The  $\text{SO}_4^{2-}$  ions may, therefore, originate from the weathering of pyrite or the leaching of other sulphides by hypothermal waters of deep origin.

The sodium chloride waters are the result of contamination by sea-water (except for springs 6 and 20) and show the possibility of sea-water intrusion at great depth (to

Table 2. Results of the chemical analyses of non-thermal karstic waters in northern Vietnam. The numbers are indicated on the map in Fig. 1.

N°	Location	T°C	TDS ( $\text{mg L}^{-1}$ )	$\text{Ca}^{++}$ ( $\text{mg L}^{-1}$ )	$\text{Mg}^{++}$ ( $\text{mg L}^{-1}$ )	$\text{Na}^+$ ( $\text{mg L}^{-1}$ )	$\Sigma \text{r}^+$ ( $\text{meq/L}$ )	$\Sigma \text{r}^-$ ( $\text{meq/L}$ )	$\text{HCO}_3^-$ ( $\text{mg L}^{-1}$ )	$\text{Cl}^-$ ( $\text{mg L}^{-1}$ )	$\text{SO}_4^{--}$ ( $\text{mg L}^{-1}$ )	Geol
24	Pho Yen	18	176.68	33.6	7.0	1.0	2.32	2.16	128.1	1.7	0.8	T
25	Son La	23	440.80	40.1	32.8	21.5	5.63	5.67	314.9	12.2	8.0	T
26	Thi Xé (HG)	24	297.73	43.5	15.5	8.6	3.82	3.81	206.9	11.5	4.1	?
27	Ha Long B.	24	414.28	89.5	17.3	15.1	5.72	5.06	249.0	24.6	8.0	C
28	Ha Long B.	24	498.63	50.2	38.0	19.8	6.65	6.78	320.0	40.2	17.0	C
29	Ha Long B.	23	350.02	73.1	2.9	18.5	4.72	4.50	207.0	30.3	9.0	C



approximately 1000 m below sea level in well 19). The great depth of the karst resulting from the subsidence of the Red River delta makes the deep flows easier (Hanoi basin, wells 18 and 19) and also the coastal karsts of Halong bay that have been partially encroached by the sea (wells 2 and 3). The proportion of sea-water in the groundwater from these wells can be estimated from the  $\text{Cl}^-$  concentration; this is  $19,000 \text{ mg l}^{-1}$  in sea-water and an average of  $20 \text{ mg l}^{-1}$  in uncontaminated karst water in the Halong region. The proportion of sea-water would thus be 42% in well 2 ( $8,842 \text{ mg l}^{-1} \text{ Cl}^-$ ) and 78% in well 3 ( $14,851 \text{ mg l}^{-1} \text{ Cl}^-$ ). However, these waters are at the same temperature ( $42^\circ\text{C}$ ) although the percentages of sea-water are very different. This indicates that there is a certain complexity in the deep flows which merits a closer study.

There is no clear relationship between the chemical facies of the waters and the temperature when the calcium sulphate waters are examined (between  $29^\circ\text{C}$  and  $68^\circ\text{C}$ ) (Fig. 3). However, the average temperature of the calcium bicarbonate waters is close to  $35.7^\circ\text{C}$  whereas that of the sodium chloride waters is  $46.7^\circ\text{C}$ . This indicates that the calcium bicarbonate waters flow at a shallower level than the sodium chloride waters or result from the mixing with small proportions of thermal water. Different chemical geothermometers have been used successfully in many studies (Michard *et al.*, 1981; Kharaka and Mariner, 1989) and are often very helpful in interpreting such geochemical processes. However, concerning thermal waters emerging from sedimentary reservoirs, the processes affecting the deep mineralized fluids prevent the use of a chemical geothermometer (Pauwels *et al.*, 1997). In the case of the thermal waters of northern Vietnam emerging from carbonate aquifers, the maximum temperatures calculated are not significant or should be considered with caution. In light of the major element analyses, the thermal waters examined in the karstic aquifers of northern Vietnam have flowed mainly through carbonate rocks. Among the 23 samples analysed, only one has a composition that indicates deep flow in contact with a crystalline basement.

With this exception, the two main features of these thermal waters are the existence of a chemical facies rich in  $\text{SO}_4^{2-}$  ions probably formed through sulphide minerals and high Na-Cl concentrations resulting from sea-water intrusion. The saline waters also result from the burying of the karstic aquifers during tectonic subsidence.

The data provided from the study of the major ions analysed in thermal karstic waters are not in themselves sufficient to detect a possible flow in deep crystalline basement, especially when the input is minor. Nevertheless, considerable chemical differences exist in thermo-mineral groundwaters from carbonate karstic formations in northern Vietnam. Also, the thermal springs studied occur neither in the same geographical context nor in the same geological units. Different hydrogeological conditions are the main factor affecting the chemical composition of these waters

and the study of the major ions has provided some reliable conclusions.

## Acknowledgements

This work was carried out within the framework of a programme of scientific co-operation between the University of Sciences, Montpellier, France, the Vietnam National Centre for Sciences and Technology (CNST), Hanoi, and the University of Mining and Geology, Hanoi, in an international programme of scientific cooperation (PICS-Vietnam) of the Centre National de la Recherche Scientifique, Paris, France and the CNST of Vietnam.

## References

- Allen, C.R., Gillespie, A.R., Han, Y., Sieh, K.E., Zhang, B. and Zhu, C., 1984. Red River and associated faults, Yunnan province, China: Quaternary, geology, slip rates and seismic hazard. *Geol. Soc. Amer. Bull.*, **95**, 686–700.
- Andreasen, D.C. and Fleck, W.B., 1997. Use of bromide chloride ratio to differentiate potential sources of chloride in a shallow, unconfined aquifer affected by brackish-water intrusion. *Hydrogeol. J.*, **5**, 17–26.
- Back, W. and Hanshaw, B.B., 1965. Chemical geohydrology. *Advan. Hydrosci.*, **1**, 49–109.
- Blondel, F., 1929. Les Phénomènes karstiques en Indochine française. *Bull. Serv. Géol. de l'Indochine*, **XVII**, fasc. 4, 1–8.
- Can, N. and Hoe, N.D., 1969. Recent faulting in coastal zone of Vietnam and its relation to seismic activities. *J. Sci. Natur. Sci., Vietnam Nat. Univ.*, **XI**, 51–56.
- Domenico, P.A. and Schwartz, F.W., 1990. *Physical and Chemical Hydrogeology*. Wiley, Chichester, UK, 824 p.
- Fénart, P., Cat, N.N., Drogue, C., Canh, D.V. and Pistre, S., 1999. Influence of tectonics and neotectonics on the morphogenesis of the karst of Halong Bay, Vietnam. *Geodin. Acta*, **12**, 193–200.
- Fisher, R.S. and Mullican, W.F., 1997. Hydrochemical evolution of sodium sulfate and sodium chloride groundwaters beneath the northern Chihuahuan desert, Trans-Pecos, Texas, USA. *Hydrogeol. J.*, **5**, 4–16.
- Fontaine, H., 1969. *Sources thermales et minérales du sud Vietnam*. Report of the 22 nd session (Mineral and thermal waters of the world; B: Overseas countries); XXIII Intern. Geol. Congress, Prague (Czechoslovakia), **19**, 63–68.
- Fromaget, J., 1941. L'Indochine française. Sa structure géologique, ses roches, ses minéralisations et leurs relations possibles avec la tectonique. *Bull. Serv. Géol. de l'Indochine*, **XXVI**, fasc. 2, 61 p.
- Geological Map of Vietnam, 1986. Scale 1/1 500 000, General Dpt. of Geology, Hanoi.
- Glazek, J., 1966. On the karst phenomena in north Vietnam. *Bull. Acad. Polonaise Sci.*, **XIV**, 45–51.
- Grimaud, D., Huang, S., Michard, G. and Zheng, K., 1985. Chemical study of geothermal waters of central Tibet (China). *Geothermics*, **14**, 35–48.
- Harrison, T.M., Leloup, P.H., Ryerson, F.J., Tapponnier, P., Lacassin, R. and Chen, Wenji, 1996. Diachronous initiation of transtension along Ailao Shan-Red River shear zone, Yunnan and Vietnam. In: *The tectonic evolution of Asia*. Cambridge University Press, 208–226.
- Hem, J.D., 1970. Study and interpretation of chemical characteristics of natural water. *U.S. Geol. Surv., Water Supply Paper*, **1473**, 363 p.
- Huang, Chu-Ching, 1978. An outline of the tectonic characteristics of China. *Eclogae Helv.*, **72**, 611–635.
- Hissou, Y., Mudry, J., Mania, J., Bouchaou, L. and Chauve, P., 1999. Utilisation du rapport Br/Cl pour déterminer l'origine de

- la salinité des eaux souterraines: exemple de la plaine du Souss (Maroc). *C. R. Acad. Sci. Paris*, 328, 381–386.
- Kharaka, Y. and Mariner, R.H., 1989. Chemical geothermometers and their application to formation waters from sedimentary basins. In *Thermal History of Sedimentary Basins*, N.D. Naeser and T. McCulloch (eds). Springer Verlag, 99–117.
- Leloup, P.H., Lacassin, R., Tapponier, P., Shärer, U., Zhong, D., Lui, X., Zhan, L. and Phan, T.T., 1995. The Ailao Shan-Red River shear zone (Yunnan, China). Tertiary transform boundary of Indochina. *Tectonophysics*, 251–285.
- Michard, G., Fouillac, C., Grimaud, D. and Denis, J., 1981. Une méthode globale d'estimation des températures des réservoirs alimentant les sources thermales. Exemple du Massif Central Français. *Geochim. Cosmochim. Acta*, 45, 1199–1207.
- Morell, I., Médina, J., Pulido, A. and Fernandez-Rubio, R., 1986. The use of bromide and strontium ions as indicators of marine intrusion in the aquifer of Torreblanca (Castellon, Spain). *Proc. 9th Salt water intrusion meeting*, Delft, 12–16 May 1986, 629–640.
- Pauwels, H., Fouillac, C., Goff, E. and Vuataz, F., 1997. The isotopics and chemical composition of CO<sub>2</sub>-rich thermal waters in the Mont-Dore region (Massif Central, France). *Appl. Geochem.*, 2, 411–427.
- Pham, V.N., Boyer, D., Van, C. and Thi, K., 1995. Propriétés électriques et structures profondes de la zone de failles du Fleuve Rouge au nord Vietnam, d'après les résultats de sondages magnéto-telluriques. *C.R. Acad. Sc., Paris*, 320, 181–187.
- Piper, A.M., 1944. A graphic procedure in the geochemical interpretation of water analyses. *Trans. Am. Geophys. Union*, 25, Hydro. paper 6, 914–927.
- Rangin, C., Klein, M., Roques, D., Le Pichon, X. and Le Van, Trong., 1995. The Red River fault system of the Gulf of Tonkin, Viet-Nam. *Tectonophysics*, 243, 209–222.
- Schoeller, H., 1962. *Les eaux souterraines*. Masson éd., Paris, 642 p.
- Shen, Zian, Tian, Honjun and Luo, Zhetan, 1989. Hydrochemical characteristics of Chinese thermomineral waters. *Steir. Beitr. Z. Hydrogeologie*, 40, 151–164.
- Standard Methods*, 1974. ALPHA, AWWA, WPCF, 13th ed., Amer. Publ. Health Assoc, 874 p.
- Tran Cau, N., 1984. *Géographie du Vietnam*. Xunhasaba éd. Hanoi, Vietnam, 241 p.
- Waring, G.A., 1965. Thermal springs of the United States and other countries of the world. A summary. *Prof. Paper Geol. Surv.* 492, 187–188.
- Xuyen, D.N., 1991. Seismic zoning of the territory of Vietnam. *Proc. 2nd Conf. on Geol. Indochina*, Hanoi, 39–44.
- Yem, N.T., 1986. Recent geodynamics of Red River Fault System. *Proc. Cong. Geol. Intern., Ho Chi Minh City*, 5–7 Dec., 405–408.